Additional Mammographic Views: A Comprehensive Guide

Louise C. Miller, R.T.(R)(M)(ARRT), CRT(M), FSBI, FNCBC Director of Education, Mammography Educators

MAMMOGR RPHY

© 2025 Mammography Educators

Additional Views Lexicon

- RL Rolled lateral
- RM Rolled medial
- RS Rolled superior
- RI Rolled inferior
- FB Caudocranial



Additional Views Lexicon – Labeling Codes

- XCCL Exaggerated craniocaudal lateral
- CV Cleavage
- ML Mediolateral
- LM Lateromedial
- AT Axillary tail
- TAN Tangential




2

Additional Views Lexicon

- SIO Superior lateral to inferior medial oblique
- LMO Lateromedial oblique
- M Magnification
- ID Implant displaced
- No label: Spot Compression





Labeling Codes (Lexicon)

The name is view (labeling code) is always preceded by identification of laterality:

• LXCCL or RXCCL



The Most Commonly Used Additional Views

- XCCL
- CV
- · LM/ML



Why do we do additional views?

- To show a specific component of the anatomy not seen on standard views.
- To provide localization of an area of concern medial/lateral or superior/inferior to the nipple



Or...

- To show an area of concern in better details
- To counteract superimposition of structures
- To triangulate a lesion



9

Commonly used additional views are done to show a specific component of the anatomy not seen on standard views.



10


Ask and Answer

- Which part of the breast do I want to visualize?
- In which projection?
- Which view will accomplish this?



Imaging the Breast in a Transverse or Axial Plane

11 12



Use of the XCCL

- Should be used on a <u>baseline</u> exam when lateral posterior breast tissue is missing on the CC view
- If glandular breast tissue on subsequent screening views is visualized back to the retromammary fat space on the MLO, an XCCL is not needed




14

XCCL required for baseline, but not on subsequent screenings **МАММОGR**


Use of the XCCL


- Should be performed on less than 10% of all patients
- Performed at 0-degree angulation
- Patient's body should be at 45-degree angle to IR
- Nipple should be pointing towards the upper corner of the image receptor

















Sternalis Muscle


- Flame-like appearance (similar to an appendix)
- Present in only 7-10% of the population
- Seen medially on a mammogram
- Often misdiagnosed as the insertion of the pectoralis muscle







23 24

6/6/25









AT – Axillary Tail

- Angle tube to axillary tail
- Approximately 30 degrees
- It is never used to localize a lesion



29

AT – Axillary Tail

It will not give you true lateral/medial or true superior/inferior orientation to the nipple



31

AT – Axillary Tail

- The AT View is used only for focal compression of the axillary tail
- Anterior to posterior orientation and compression




30

90-degree (True) Lateral

- LM Lateromedial
- ML Mediolateral





Use of the Lateral

- Shows effects of gravity on air fluid levels (Milk of Calcium)
- Used as a "tie breaker" view (to overcome superimposition of structure)
- Visualizes the breast in the sagittal plane (demonstrates an area of concern superior or inferior to the nipple)



EDUCA

34

• When you did the MLO, you showed the lateral breast in better detail; The LM shows the medial breast in better detail

Why do the LM?

• The LM takes advantage of the lateral mobile border of the breast and thus facilitates positioning



Why do the LM?

- The posterior medial breast is hardest part of the breast to image and the area most often missed on the MLO
- If done properly, by off-setting the IR into the contralateral breast, you will be able to go deeper against the chest wall



35

Why do the LM?

• There is no issue of the contralateral breast impeding the path of the compression paddle




2000 0 1 1 0

37

Improperly positioned LM with breasts separated, so the middle of the IR is centered on midsternal line. This excludes deep medial breast tissue on the side you are imaging.

38

Properly positioned LM with breasts separated so the *top* edge of the IR is centered on midsternal line and the width of the IR pressing against the contralateral breast.



MAMMOGR&PH

39

ML - Mediolateral

The opposite breast must be pulled back to allow the compression paddle to pass and may therefore eliminate visualization of deep medial breast tissue.





40

Additional Views to Facilitate Imaging of Difficult Body Habitus

- LMO Lateromedial Oblique
- FB Caudocranial (From Below)



41

LMO – Lateromedial Oblique

- Degree of angulation should be 90 degrees plus or minus the standard degree of angulation.
- Average 45-degree RMLO would be 135 degrees for RLMO



43 44

LMO and FB Views

- Used when a standard MLO or CC is impossible
- Kyphotic patients
- Males with prominent pectoral muscles



42







MAMMOGR RPHY

FB - Caudocranial

- Machine is turned 180 degrees opposite the CC
- Direction of the beam is Caudal to Cranial



45

Additional Views for Clarification of AOC

- TAN
- Spot compression
- Spot compression with MAG
- Rolled views



48 47



FB

CC


To prove the existence of dermal calcificationsEnhanced visualization of palpable masses that may otherwise be superimposed on glandular breast tissue




MAMMOGR RPHY

46

6/6/25





49 50





51 52


Localization for Verification of Skin Calcifications

- Decreased with use of DBT
- Set up as the same as needle localization
- Determine which quadrant the calcifications are located





E D U C A T O R

53




54








55

6/6/25
























6/6/25











Remember...

- You must stimulate compression when making measurements on the breast
- Mark the center of the target area with a surgical marker so you can make appropriate corrections on subsequent images, if needed



70

69

Remember...

- Rolled views used to overcome superimposition of structures by changing the orientation of the beam to the breast
- Lateral views used to overcome superimposition of structures by changing the orientation of the breast to the beam





71 72


Rolled Views


- RCCRM Right CC superior breast rolled medial RCCRL Right CC superior breast rolled lateral














Imaging of Augmented Breasts

Full implant views:

- Should be done with only enough compression to immobilize the breast to prevent motion unsharpness
- Curved paddle can be used (if available)
- Appropriate technique (usually manual) should be used



79 80

Imaging of Augmented Breasts ID views: • Depending on implant mobility, can be performed with taut compression • Half paddle can be used for patients with small amount of natural breast tissue • Appropriate techniques (used for patients without implants) should be used • Patient can be positioned from behind (with tech standing and/or patient seated)


Half Paddle


MAMMOGR&PHY

Image Courtesy of Robyn Hadley, R.T.(R)(M)


81


MAMMOGR & PHY





83





Conclusion

Additional views:

- Are helpful in identifying true location of areas of concern
- Are used for diagnostic workups
- Can provide valuable information to aid in diagnosis of breast cancer



87

References

Miller, L. C., RTRM. (2015). Mammography Positioning Guidebook. CA: Mammography Educators.

Miller, L. C., R.R.(R)(M)(ARRT), Lehmann, T. L., B.S. R.T. (R)(M)(ARRT). (2020). Image Quality & Positioning Problem-Solving for Breast Imagers: Meeting EQUIP Standards (1st ed.). San Diego, California: Mammography Educators.

Long, S. M., Miller, L. C., Botsco, M.A., & Martin, L. L. (2010). Handbook of Mammography (5th ed.). Edmonton: Mammography Consulting Services.

Derenberger, Dawn, Hadley, Robyn., "Most Commonly Used Additional Views, Part 3: Defining Structures and Clarifying Presence of Abnormalties." SBI News Issue 1,



88

