

The Importance of Standardized Positioning:

The Basics of Mammography That You Were Never Taught

Louise C. Miller, R.T.(R)(M)(ARRT), CRT(M), FSBI, FNCBC
Director of Education, Mammography Educators



MAMMOGRAPHY
EDUCATORS

© 2025 Mammography Educators

1

The screenshot shows the FDA website with the article title "Poor Positioning Responsible For Most Clinical Image Deficiencies, Failures". The article text states: "Mammography combines 'the science of imaging and the art of positioning' [1]. Although there have been many significant and exciting changes to the technology of mammography since the passage of MQSA in 1999, including the introduction of full-field digital mammography (FFDM) and digital breast tomosynthesis (DBT), one aspect of mammography that remains unchanged and critically important is proper patient positioning. Positioning is so important because only those portions of the breast which are included on the mammograph image can be evaluated for signs of cancer. Any portion of the breast which is not imaged cannot be evaluated, and cancers in those portions of the breast can be missed. In a 2002 study, the '[s]ensitivity [of mammography] dropped from 84.4% among cases with passing positioning to 66.3% among cases with failed positioning' [2]. Poor positioning has been found to be the cause of most clinical image deficiencies and most failures of accreditation. In 2015, the American College of Radiology (ACR), the largest FDA-approved accreditation body (AB), found that of all clinical images which were deficient on the first attempt at accreditation, 92% were deficient in positioning. Also, in ACR-accredited facilities, 79% of all unit accreditation failures in 2015 were due to

2

US Food and Drug Administration

- "In a 2002 study, the '[s]ensitivity [of mammography] dropped from 84.4% among cases with passing positioning to 66.3% among cases with failed positioning'."
- "Poor positioning has been found to be the cause of most clinical image deficiencies and most failures of accreditation." (92%)



MAMMOGRAPHY
EDUCATORS

3

Decreased Sensitivity

- 84.4% with proper positioning
 - 66.3% with failed positioning
- = 18.1% decreased sensitivity**



MAMMOGRAPHY
EDUCATORS

4

Quality Standards – Why?

ALL industries have established *standardized* methods in the performance of tasks to:

- Establish and maintain quality
- Increase consumer satisfaction
- Increase profit
- Reduce possibility of litigation
- Reduce errors



MAMMOGRAPHY
EDUCATORS

5


How do we reduce medical errors?

- Standardization
- Consistency
- Reproducibility



MAMMOGRAPHY
EDUCATORS

6



MAMMOGRAPHY
EDUCATORS

7

Standardized Technologist Training for General Radiology



MAMMOGRAPHY
EDUCATORS

8

Standardization

We all:

- Position the same way for every body part
- Position in the same sequence
- Set up the machine before we bring in the patient
- Position the whole patient, not just the body part



MAMMOGRAPHY
EDUCATORS

9


In General Radiology...

All training is competency based and a technologist's skills will be evaluated for *positioning techniques*, as well as *clinical image evaluation*.



MAMMOGRAPHY
EDUCATORS

10



MAMMOGRAPHY
EDUCATORS

11

Consistency and Ergonomics

- **M**achine
- **P**atient
- **B**ody Part (Breast)



MAMMOGRAPHY
EDUCATORS

12

We position the **whole patient**,
not just the body part!



MAMMOGRAPHY
EDUCATORS

13



MAMMOGRAPHY
EDUCATORS

14

All exams are done using the *same*
positioning technique, in the *same* sequence.



MAMMOGRAPHY
EDUCATORS

15

Why???

- Consistency
- Reproducibility
- Efficiency
- Proficiency
- Use of proper body mechanics



MAMMOGRAPHY
EDUCATORS

16

But in mammography... We are "all over the map"

- LCC, LMLO, RMLO, RCC
- RCC, LCC, RMLO, LMLO
- RMLO, RCC, LMLO, LCC
- LCC, RCC, LMLO, RMLO
- RCC, RMLO, LMLO, LCC
- LCC, LMLO, RCC, RMLO
- LMLO, LCC, RCC, RMLO



MAMMOGRAPHY
EDUCATORS

17

My Suggestion

- CCs first
- Then do the MLO on the side you just finished the CC
- Finish with the other MLO

Example: RCC, LCC, LMLO, RMLO



MAMMOGRAPHY
EDUCATORS

18

So why is this true for all body parts in
radiology **EXCEPT** in Mammography???



MAMMOGRAPHY
EDUCATORS

19

In Mammography...


- Most technologists have not been taught a standardized method of positioning
- Most technologists have been taught various positioning methods by various technologists



MAMMOGRAPHY
EDUCATORS

20

Most technologists *do not* practice a standardized method of positioning or position the same way.



MAMMOGRAPHY
EDUCATORS

21

No Standards for Mammography Positioning

There are standards for *WHAT* images should look like, but not *HOW* you get to that point!



MAMMOGRAPHY
EDUCATORS

22

So the problem is...

No standardization or follow-through, which means:

- Less consistency and reproducibility
- More repeats and rejects
- More accreditation failures
- Increased radiation exposure
- MISSED BREAST CANCERS



MAMMOGRAPHY
EDUCATORS

23

How can we make things better?

Consistency and Reproducibility



MAMMOGRAPHY
EDUCATORS

24

Mammography Positioning Techniques

Should be:

- Consistent
- Reproducible
- Efficient
- Proficient
- Based on sound ergonomic principles



MAMMOGRAPHY
EDUCATORS

25

Common Work-Related Injuries

- Wrists
- Shoulders
- Back
- Knees
- Hips



MAMMOGRAPHY
EDUCATORS

26

YES!!!



NO!!!



MAMMOGRAPHY
EDUCATORS

27

YES!!!




NO!!!



MAMMOGRAPHY
EDUCATORS

28



29

Do Standardized Positioning Techniques for Mammography Work?

- Scientific studies prove that standardized positioning techniques improve quality
 - Visualization of more posterior and lateral breast tissue, IMF and pectoralis muscle on the CC
- Unpublished data showed that standardized positioning techniques decrease repeats, rejects and technical callbacks by up to 50%!

30

Mammography Positioning Standards in the Digital Era: Is the Status Quo Acceptable?

Ashley L. Huppe¹
Kelly L. Overman²
Jason B. Gatewood³
Jacqueline D. Hill⁴
Louisa C. Miller⁵
Marc F. Inciardi¹

OBJECTIVE. The objective of our study was to evaluate positioning of full-field digital mammography (FFDM) and digital breast tomosynthesis (DBT) compared with film-screen (FS) mammography positioning standards.


MATERIALS AND METHODS. A retrospective study was conducted of consecutive patients who underwent screening FFDM in 2010–2012 and DBT in 2012–2013 at an academic institution. Examinations were performed by five experienced technologists who underwent updated standardized positioning training. Positioning criteria were assessed by consensus reads among three breast radiologists and compared with FS mammography data from a 1993 study by Bassett and colleagues.

RESULTS. One hundred seventy patients (n = 340 examinations) were analyzed, showing significant differences between FFDM and DBT examinations (p < 0.05) for medial or inferior skin folds (FFDM vs DBT: craniocaudal [CC] view, 16% [n = 56] vs 23% [n = 77]; mediolateral oblique [MLO] view, 35% [n = 118] vs 45% [n = 154]); inclusion of lateral glandular tissue on CC view (FFDM vs DBT, 73% [n = 247] vs 81% [n = 276]), and concave pectoralis muscle shape (FFDM vs DBT, 36% [n = 121] vs 28% [n = 95]). In comparison with Bassett et al. data, all positioning criteria for both FFDM and DBT examinations were significantly different (p < 0.05). The largest differences were found in visualization of the pectoralis muscle on CC views and the inframammary fold on MLO views, inclusion of posterior or lateral glandular tissue, and inclusion of skin folds, with DBT and FFDM more frequently exhibiting all criteria than originally reported Bassett et al. findings.

CONCLUSION. DBT and FFDM mammograms more frequently include posterior or lateral tissue, the inframammary fold on MLO views, the pectoralis muscle on CC views, and skin folds than FS mammograms. Inclusion of more breast tissue with newer technologies suggests traditional positioning standards, in conjunction with updated standardized positioning training, are still applicable, at the expense of including more skin folds.

Source: American Journal of Roentgenology; 209, December 2017

31



32

Reasonable Expectations for the CC

	Positioning Criteria	FFDM	DBT	Bassett
CC View	Pec Muscle Visualized	48%	50%	32%
	No Motion	100%	98%	-
	Lateral Glandular Tissue Included	73%	81%	37%
	Nipple in Profile	83%	85%	89%
	Skin or fat folds	39%	47%	10%
	Medial Location	16%	23%	-
	Lateral Location	29%	32%	-
	Visualization of Cleavage	41%	34%	-
	Requires More Than One View	5%	7%	-



33

Reasonable Expectations for the MLO

	Positioning Criteria	FFDM	DBT	Bassett
MLO View	Visualization of Pec Muscle to PNL	86%	87%	81%
	Concave Pec	36%	28%	-
	Straight Pec	41%	46%	-
	Convex Pec	23%	26%	-
	Wide Margin at Top of Pec	95%	93%	-
	No Motion	98%	97%	99%
	Posterior Glandular Tissue Included	90%	94%	77%
	Nipple in Profile	89%	92%	88%
	Skin or fat folds	53%	62%	15%
	Upper Location	25%	27%	-
	Lower Location	35%	45%	-
	Visualization of Inframammary Fold	81%	85%	49%
	Requires More Than One View	13%	17%	-



34

Reasonable Expectations

- Our patients have different and often challenging body habitus
- Their breast size, shape, mobility and tenderness are hugely variable



35

Reasonable Expectations

Even the “perfect” patient, in terms of body habitus, breast mobility, etc. may provide a challenge that inhibits the technologist’s ability to position and compress properly.



36



37



38



39

But regardless of these variables, we **CAN** improve image quality by **using standardized positioning techniques** and developing a strong knowledge-based foundation that depends on the technologist's understanding of correlative anatomy.

40

RCC: Steps 1-9 (Front)



41

Quick Steps for the CC (Left)

- ___ 1. Elevate breast/IMF (until the PNL is perpendicular to the chest wall)
- ___ 2. Adjust IR height (so top edge is parallel with elevated IMF)
- ___ 3. Pull the breast onto the IR with both hands (left hand on top; right hand on bottom) and at the same time ask the patient to step forward into the machine (no leaning) and have them turn their face towards you
- ___ 4. Switch hands, so your right hand is on the top (palm down) and anchor the breast with the base of your right thumb.
- ___ 5. Lift the opposite/contralateral breast onto the IR with your left hand, palm facing up; then ask the patient to turn their right hip forward.
- ___ 6. Guide the patient's head forward and around the face shield, if possible
- ___ 7. Place your left elbow and forearm at the mid thoracic region (where their bra clasp would be) and gently push the patient forward
- ___ 8. Relax patient's left shoulder with your left hand, if possible
- ___ 9. Pull superior breast tissue forward by placing the base/edge of your right thumb on top of the breast against the chest wall, then apply compression while continuing to "push" the patient forward



Source: Mammography Educators

42

RMLO: Steps 1-10



43

Quick Steps for the MLO (Left)

- ___ 1. Stand perpendicular to the patient with your sternum pressing against patient's right humerus
- ___ 2. Lift patient's left shoulder/arm up over the corner of the IR with your right hand in the patient's axilla. At the same time, your right hand should "meet" your left hand in the axilla and help to lift the patient's left shoulder up and over the IR
- ___ 3. IR is placed in back of axilla (just interior to latissimus dorsi)
- ___ 4. Patient's left hand should be resting on bar, with their elbow bent behind the IR
- ___ 5. Place your left hand on patient's left shoulder (if possible) to keep their shoulder relaxed and down
- ___ 6. Your right hand, with palm facing up, slides down lateral side of the breast to pull on lateral breast tissue and smooth out any skin folds
- ___ 7. Once your right hand is at the bottom of the breast, turn your hand over so that your hand is now palm down on the breast with the base of your thumb just anterior to the IMF
- ___ 8. Push the breast up and out with the base of your thumb, keeping continuous contact with the breast
- ___ 9. At the same, ask the patient to lift and flatten their other breast, as needed
- ___ 10. Continue to hold the breast in the up and out position until compression is complete



Source: Mammography Educators

44


Does it work??



MAMMOGRAPHY
EDUCATORS

45


BEFORE 13.6 cm STANDARDIZED POSITIONING + 3.0 cm AFTER 16.6 cm



MAMMOGRAPHY
EDUCATORS

46


BEFORE 14.8 cm STANDARDIZED POSITIONING + 2.0 cm AFTER 16.8 cm



MAMMOGRAPHY
EDUCATORS


47

BEFORE 13.9 cm STANDARDIZED POSITIONING + 0.7 cm AFTER 14.6 cm




MAMMOGRAPHY
EDUCATORS


48



49



50



51



52

It is ALL our responsibility to make sure that ALL patients receive the highest quality mammogram achievable.



53

Back To The Basics

Going back to the basics strengthens your foundation.

In Mind of Truth



54

Mammography should be taught according to scientific principle, **NOT ANECDOTE!**



55

Anecdote vs. Science



56

Anecdote vs. **Science**



57

SCIENCE:

ANATOMY – PHYSIOLOGY – PHYSICS



58

You lucked out...



59

Except...

- Newton's Third Law: for every action there is reaction
- Law of gravity



60

Anatomy and Physiology

- As they relate to mammography positioning, using general radiology principles...



MAMMOGRAPHY
EDUCATORS

61

Goals for **General Radiology** Positioning

- Bring the body part back to its true anatomical position OR the position that will best visualize the body part
- Use palpable and visible anatomical landmarks for positioning and clinical image evaluation
- Use consistent and reproducible methods



MAMMOGRAPHY
EDUCATORS

62

Goals for **Mammography** Positioning


Bring the breast back to its natural anatomical position (*with the nipple perpendicular to the chest wall as possible*) on both screening views to maximize visualization of breast tissue and to avoid superimposition of structures.



MAMMOGRAPHY
EDUCATORS

63

Anatomy of the Breast



MAMMOGRAPHY
EDUCATORS

64

Normal or natural position of the breast is when the nipple is perpendicular to the chest wall.



65

When positioning for mammography we need to bring the breast back to its 'normal' position.




66

In order to accomplish this and include the maximum amount of breast tissue, we must consider the anatomical landmarks that will be used for positioning and clinical image analysis.



67



68


Perimeter of the Breast



MAMMOGRAPHY
EDUCATORS

69


Perimeter Used for Positioning and Clinical Image Analysis



MAMMOGRAPHY
EDUCATORS

70

Pectoralis Used for Positioning and Clinical Image Analysis




MAMMOGRAPHY
EDUCATORS

71

PNL Used for Positioning

Elevate the breast so that the PNL is as close as possible to perpendicular to the chest wall.




MAMMOGRAPHY
EDUCATORS

72

PNL Used for Clinical Image Analysis

PNL measurement of CC should be within 1cm of the PNL measurement on the MLO.




73

The MLO

- Inclusion of all breast tissue within perimeter
- Pectoral muscle fully visualized
- Tissue well separated
- Tissue visualized back to the retromammary fat space
- IMF



74



75

The MLO

Visualization of the pectoral muscle:

- The pectoralis muscle is not really part of the breast
- However, it serves as an important anatomical landmark for positioning and image evaluation



76

The absence or presence of these characteristics will tell you exactly what you did right or wrong when positioning and therefore, whether you included or excluded breast tissue!



MAMMOGRAPHY
EDUCATORS

77

Remember

There are only two margins for error:

- 1) The way the **machine** is set up (i.e., height, angle, compression paddle size, etc.)
- 2) The way the **patient** is “set up”: both feet, hips, and shoulders facing forward



MAMMOGRAPHY
EDUCATORS

78

The MLO

- **Width:** wide margin at the top
- **Length:** down to the level of the PNL
- **Shape:** convex/straight




MAMMOGRAPHY
EDUCATORS

79

The MLO

WIDTH of the Muscle

There should be a wide margin of the pectoralis muscle at the top of the image (in the axilla).



MAMMOGRAPHY
EDUCATORS

80

The MLO

EQUIPMENT: Width of the muscle is related to the placement of the IR in the axilla.

The back corner of the IR should be placed just anterior to the latissimus dorsi.



MAMMOGRAPHY
EDUCATORS

81



MAMMOGRAPHY
EDUCATORS

82

The MLO


PATIENT: Width of the muscle is related to the position of the patient.

The patient must be turned into the machine with both feet, hips and shoulders as far forward as possible. The patient's shoulder should be down and relaxed and if possible, held in position by the technologist.



MAMMOGRAPHY
EDUCATORS


83



MAMMOGRAPHY
EDUCATORS

84

Width of the Muscle




85

The MLO

LENGTH of the Muscle

Should be visualized down to the level of the PNL.



86

The MLO


EQUIPMENT: Angle for the MLO

- Angle to the free margin of the pectoralis muscle
- Keep angulation consistent
- Steeper angle for patients with longer thoraxes and small breasts
- Lesser angles for shorter thoraxes and larger breasts




87

Proper degree of angulation



Angle too steep



88

Recommended Angulation for the MLO

- Depends on body habitus
- Maintain consistency from year to year*

*An MLO angled at 56-degrees one year will look markedly different than an MLO angled at 42-degrees the next year.



MAMMOGRAPHY
EDUCATORS

89

Keep Angles Consistent

Use 5-degree increments – no more 43, 48, 52 degrees

- 40 degrees for shorter, heavier patients with large breasts
- 45 degrees for average patients
- 50 degrees for tall, thinner patients with smaller breasts
- 35 degrees for patients who have undergone reduction



MAMMOGRAPHY
EDUCATORS

90

The MLO

PATIENT: Length of the muscle is related to the position of the patient.


The patient must be turned into the machine with both feet, hips and shoulders as far forward as possible, as not to impede the progress of the compression paddle.




MAMMOGRAPHY
EDUCATORS

91

Proper degree of angulation



Angle too steep




MAMMOGRAPHY
EDUCATORS

92

OR...

The patient is not facing the machine properly.




93

The MLO

SHAPE of the Muscle

The muscle should be convex or straight.



94

The MLO

EQUIPMENT: The shape and opacity of the muscle is related to the height of the IR.

The top of the IR should be positioned at the height of the sternoclavicular joint or halfway between the top of the shoulder and the axillary crease.



95


The MLO

PATIENT: The shape and opacity of the muscle is related to the relaxation of the pectoralis muscle.


Patient's shoulder, arm and hand must be relaxed with the elbow bent and relaxed behind the IR.




96




97




98



99



100




101



102




103




104

The “Sagging” Breast




105

The “Sagging” Breast



106

Stages of Breast Cancer Development



107


Position of the Breast

- Breast must be held in the “up and out” position to bring the breast back to its “normal” position (nipple perpendicular to the chest wall)
- Maintained by adequate compression
- Don’t let go until compression is complete




108

Solution for the “Sagging” Breast



Hold the breast in up and out position.



Compress.




MAMMOGRAPHY
EDUCATORS

109



The CC

- Include maximum amount of breast tissue in the axial/transverse plan
- Visualization of medial breast tissue
- Visualization of pectoralis muscle on approximately 50% of all CCs
- PNL within 1cm of PNL on MLO



MAMMOGRAPHY
EDUCATORS

110



MAMMOGRAPHY
EDUCATORS

111

The CC

- The CC is done to provide an “orthogonal” view from the MLO
- When added to the MLO, it increases breast cancer detection
- While the CC is not primarily done to visualize medial breast tissue (cleavage), it is a “by-product” of a good CC
- If you miss breast tissue on the MLO, it will be medial tissue, so it’s important to try to include that on the CC view



MAMMOGRAPHY
EDUCATORS

112

The CC

Is it the equipment or the patient?

Equipment:

- IR too high or too low
- Compression paddle size

Patient:

- Facing towards the machine with both feet, hips and shoulders forward



MAMMOGRAPHY
EDUCATORS

113

Due to lack of anatomical landmarks,
positioning techniques are extremely important!!



MAMMOGRAPHY
EDUCATORS

114

The CC

Common problems:


- Short CC (PNL measurement on CC is less than 1 cm of PNL measurement on the MLO)
- No pec muscle (remember the 50% rule!)
- No visualization of deep medial breast tissue (cleavage)

All of these are most often related to positioning
technique vs. equipment and patient position.
(Although they are still important!)



MAMMOGRAPHY
EDUCATORS


115



MAMMOGRAPHY
EDUCATORS

116

Meet, Beat, Repeat



MAMMOGRAPHY
EDUCATORS

117

Focus On...

- Consistency
- Reproducibility
- Efficiency
- Proficiency
- Ergonomic principles



MAMMOGRAPHY
EDUCATORS

118

Mammography Saves Lives!

But it is up to you.....

Even the best radiologist, in the best breast center cannot diagnose a cancer that is not included on the image.



MAMMOGRAPHY
EDUCATORS

119



MAMMOGRAPHY
EDUCATORS

120

References

"Poor Positioning Responsible For Most Clinical Image Deficiencies, Failures." U.S. Food and Drug Administration, 29 November 2017.
https://public4.cvsfreszer.com/browser/FDA/77_12-2022108-32/items/www.fda.gov/radiation-emitting-products/mma-insists/poor-positioning-responsible-most-clinical-image-deficiencies-failures

Miller, Louise C. "A Technologist's Perspective of the FDA Report 'Poor Positioning for Most Clinical Image Deficiencies, Failures.'" SBI News, 3 June 2016.
<https://mammographyeducation.com/a-technologists-perspective-of-the-fda-report/>

Miller, Louise C. "What Every Mammography Technologist Would Like Their Radiologist to Know About: Our Patients." SBI Newsletter, 2015.
<https://mammographyeducation.com/what-every-mammography-technologist-would-like-their-radiologist-to-know-about-the-role-of-the-technologist/>

James, John T. PhD. "A New, Evidence-based Estimate of Patient Harms Associated with Hospital Care." Journal of Patient Safety, September 2013.
https://journals.lww.com/journalofpatientsafety/Fall2013/09000/A_New_Evidence_based_Estimate_of_Patient_Harms?stop



Thank You!

Services we offer, include:

- Onsite Positioning Training
- Assistance with Accreditation & Inspection
- Live Webinars and Conferences
- On-Demand Continuing Education

For questions or more information:

619-663-8269

mammographyeducators.com

info@mammographyeducators.com

