

*An accolade for Ms. Louise C. Miller's life-time achievement*






Louise C. Miller, R.T.(R)(M)(ARRT),  
CRT(M), FSBI, FNCBC



László Tabár, MD, FACR (Hon)

1

*László Tabár, MD  
Tibor Tot, MD, Peter B. Dean, MD*




Understanding the Breast  
in Health and Disease

2



3


**Narrated book:** <https://www.dropbox.com/sc/fi/fu6d9j178mjaftewsbns/PARTS-I-IV-MERGED.mp4.patched.mp4?rlkey=ohw509twformiz52pfh4enf0r&dl=0>

**ENGLISH**

*László Tabár, MD  
Tibor Tot, MD, Peter B. Dean, MD*




Understanding the Breast  
in Health and Disease





The book page by page: <https://www.dropbox.com/sc/fi/alp5pxw3i0pvdxojsburq/Vol-I-High-qual-FINALx.pdf?rlkey=pj3pvif3ldn1pyc7gg2fxo0b1&dl=0>

4

Narrated book in Spanish PARTs I – IV (see next slide)

*Dr. László Tábar,  
Dr. Tibor Tot, Dr. Peter B. Dean*



Bases para entender la mama en la salud y en la enfermedad

En 3D

The book in Spanish, page by page: <https://www.dropbox.com/scl/fi/aldjrsa2tck2co16mymkf/Vol-I-SPANISH.pdf?rlkey=5jb44xbi9215nm8ucl5yvvxqg&dl=0>

5

**SPANISH**

**PART I**

<https://www.dropbox.com/scl/fi/26dfw8fb0xr1hq2gk5fpt/3D-Spanish-Part-I.mp4.patched.mp4?rlkey=mzjvmu9qkcth3jiwuc85xyvgs&dl=0>

**PART II**

<https://www.dropbox.com/scl/fi/jo9iy8aroilrh43wwzdf/3D-Spanish-Part-II.mp4.patched.mp4?rlkey=9dn2oyzmbzk3q4tev79oj6hif&dl=0>


**PART III**

<https://www.dropbox.com/scl/fi/gt0gwj73xq2n5ltjmp6bk/3D-Spanish-Part-III.mp4.patched.mp4?rlkey=by30tw2tj9441453zku47i1st&dl=0>


**PART IV**

<https://www.dropbox.com/scl/fi/o1s8jp050jn0l2jzgl9xi/3D-Spanish-Part-IV.mp4.patched.mp4?rlkey=tq5afmd769cexlp1td5vzp6vg&dl=0>

6



7



8

It is our firm belief that competence in breast image interpretation can be best achieved by correlating the large format 3D histopathology of the normal and abnormal breast tissue with

- Mammograms
- Ultrasound, MRI and
- Other imaging methods.



9



The magical 3-D, subgross, thick section histology

10



Plastinated, large format subgross histopathology specimen as teaching tool.

11



The magical 3-D, subgross, thick section histology

12




13



The magical 3-D, subgross, thick section histology

14




The magical 3-D, subgross, thick section histology

15



The magical 3-D, subgross, thick section histology

16



The magical 3-D, subgross, thick section histology

17



The magical 3-D, subgross, thick section histology

18


Progress in histologic-mammographic correlation can only be made by examining a histologic specimen of *greater length, width and depth.*  
*Tot, T, L. Tabar, P.B. Dean*  
*Virchows Arch (2000) 437:338-344.*

19


*12-18 lobes*

A drawing of a surgical preparation of a breast by Sir Astley Paston Cooper, who injected wax of different colors into the end of each duct in the nipple to outline the individual lobes.


20




21




22




23




24




25




26




27




28



29




30



Understanding anatomy, physiology


31




1. Normal fibroglandular tissue (unequivocal)

The most frequent finding at mammography


32




33




34




35




36




37




38




39




40




41



42



43




44


Breast MRI – 3D histologic correlation of normal breast tissue




45



46




47



Structures analogous to these 3D images of the breast can also be found in the plant world.

48



49

The four basic structural elements (lobules, milk ducts, fat and fibrous connective tissue) of normal breast tissue are always present in every woman's breast, even if their relative proportions may vary widely. This variation results in five characteristic presentations on the mammogram, called the mammographic parenchymal patterns.

- These patterns can affect the mammographic detectability of a small breast cancer.
- Our classification is a "qualitative classification" based on large format histopathologic-mammographic correlation, taking into account the relative proportion of the four basic building blocks of the breast.

50

lobules, milk ducts, fat and fibrous connective tissue

mammographic detectability of a small breast cancer.


51

This classification is a "qualitative classification" based on large format histopathologic-mammographic correlation. A thorough comparison between the large format histologic image of the surgically removed breast tissue and the mammograms resulted in this classification, which is based on the relative proportion of the four basic building blocks of the breast.

The American BIRADS classification is a simple "quantitative" classification, based on the amount of radiopaque/dense tissue seen on the mammogram: fatty replaced breast, <25% dense tissue, 50% dense breast tissue, >75% dense breast tissue. The BIRADS classification does not take the underlying histologic composition into account.

52

**Pattern I.**




Approximate relative proportion of the four building blocks

Building Block	Relative Proportion
Nodular density	~27%
Linear density	~13%
Homogeneous fibrous tissue	~6%
Radiolucent adipose tissue	~54%


 Nodular density  
 Linear density  
 Homogeneous fibrous tissue  
 Radiolucent adipose tissue

53

BI-RADS **A**      BI-RADS **B**



*Pattern II*      *Pattern III*



The dynamic change of Pattern I over time

Age Group	Pattern I (%)
40-49	27%
50-59	13%
60-69	6%
70+	6%


Combined Pattern II and Pattern III

Age Group	Combined Pattern II and III (%)
40-49	73%
50-59	87%
60-69	94%
70+	94%

Tabar L, T Tot & PB Dean *The Art and Science of Early Detection with Mammography*, Thieme Verlag


54

**Pattern I.**




55


Cooper's "ligament"




56




57




58




59




60




61




62




63




64




65



66



67



68

BI-RADS D

Pattern IV

Pattern V

Age Group	Prevalence
40 - 49	11%
50 - 59	12%
60 - 69	12%
70+	13%

Age Group	Prevalence
40 - 49	16%
50 - 59	3%
60 - 69	2%
70+	2%

69

Simple adenosis, Pattern IV

70


Adenosis pattern: hyperplasia and hypertrophy of the TDLU.

71

Pattern IV - adenosis pattern

Age Group	Prevalence
40 - 49	11%
50 - 59	12%
60 - 69	12%
70+	13%

72



73

© International Epidemiological Association 2000 Printed in Great Britain                      International Journal of Epidemiology 2000;29:11-19


### Mammographic parenchymal patterns and risk of breast cancer at and after a prevalence screen in Singaporean women

RW Jakes,<sup>a</sup> SW Duffy,<sup>b</sup> FC Ng,<sup>c</sup> F Gao<sup>a,d</sup> and EH Ng<sup>c</sup>


<b>Background</b>	The objective of this study was to assess the effect of mammographic parenchymal patterns on risk of breast cancer detected at first screen or in the period following a negative screen.
<b>Methods</b>	The study utilizes a nested case-control design with 132 breast cancer patients detected at first screen (from a total of 29 193 screened) and 42 breast cancer patients detected in the period following the first screen. These patients were matched to 348 screened-negative controls. The mammograms were classified according to Tabar's classification for parenchymal pattern and statistical analysis was done by conditional logistic regression.
<b>Results</b>	The risk of breast cancer for women with Tabar pattern IV was significantly high when compared to the remaining patterns (odds ratio 2.59). Risk factors for Tabar pattern IV coincided largely with established risk factors for breast cancer.
<b>Conclusion</b>	The study confirms the increased risk of breast cancer associated with Tabar pattern IV (approximately Wolfe pattern F2), in an Asian population. This pattern is associated with nullparity and high educational status and is strongly associated with grade 3 cancers.
<b>Keywords</b>	Breast cancer, mammography, parenchymal patterns, Asian women, case control study
<b>Accepted</b>	27 July 1999

Downloaded from <http://ije.oxfordjournals.org/> by guest on


74



75




76



77



78



79

*Demonstration of the problem*

Invasive breast cancer(s) may be hidden in the dense fibrous tissue

*Example 02*

80

**Breast MRI**

Comparison of the ABUS slice 8/18, lateral compression, (a) with breast MRI images.  
**Description of the MRI findings:** there is a rounded 15x14x19 mm solitary tumor mass with irregular borders in the right breast, 8 cm deep to the nipple, in the direction of 9 o'clock (b,c,e,f.). The lesion has rim enhancement with rapid wash-in and washout in delayed phase. Both morphology and kinetics are characteristic for a **malignant tumor**. A 15 mm pathological lymph node is seen in the right axilla (g).

81

**Imaging the dense portion of the breast: Description of the problem and a promising solution.**

**Multimodality approach to the detection of breast cancer: FFDM combined with **Sonographic CT** or **3D Sonogram** (R. Godette) ??? New term? Ideas?**

82


**Automated breast ultrasound**

**ABUS**

83

*Sequence of events*

84




85




86



87



88




89




90




91




92




93




94




95




96




97




98



99



100



101

Recommended literature

**Breast Cancer: The Art and Science of Early Detection with Mammography**


www.thieme.com

102


### Where to look?

- *To detect AABs*
- *To detect DABs*
- *To detect diffusely infiltrating breast malignancy of mesenchymal origin*


103




104




105



106



107





108

*Side by side, step by step viewing*  
*Areas that require special attention*

Locations with high frequency of abnormal lesions ("forbidden areas"):


- "Milky way"
- The medial half of the breast

Circular    Stellate    Nonspec. asymm. density

109

*Side by side, step by step viewing*  
*Areas that require special attention*



61.2%	17.0%
1.1%	11.1%
9.6%	

The medial half of the breast

110


*Side by side, step by step viewing*  
*Areas that require special attention*

Locations with high frequency of abnormal lesions ("forbidden areas"):

- "Milky way"
- The medial half of the breast
- Retroglandular clear space ("no man's land")

111

*Side by side, step by step viewing*  
*Areas that require special attention*



The retroglandular clear space ("no man's land")

112


*Side by side, step by step viewing*  
*Areas that require special attention*

Locations with high frequency of abnormal lesions  
 ("forbidden areas"):

- "Milky way"
- The medial half of the breast
- Retroglandular clear space ("no man's land")
- The retroareolar area


113

*Side by side, step by step viewing*  
*Areas that require special attention*



114


*Aim of perception*  
 Disruption of the harmonic structure



Parenchymal contour change

115



*Aim of perception*  
 Disruption of the harmonic structure



Parenchymal contour  
 retraction (*tent sign*)


116

*Aim of perception*  
**Disruption of the harmonic structure**





Parenchymal contour retraction (**tent sign**)

117




118



119



120




121

53-year-old asymptomatic woman, called back from mammography screening for assessment of the finding on her mammogram.


Case 1289

From the Departments of Mammography  
Surgery and Clinical Pathology  
Falun Central Hospital, Sweden ©


122




123




124




125




126




127



128



129



130

*Histopathology*

12x10 mm solitary invasive intralobular BCNO. pN 0/2

131

*Follow-up*

Nine years and seven months after diagnosis and treatment, the patient is well with no signs of recurrence.


132

*Breast imagers' expectations for help from AI when reading mammograms*




I am truly enthused by the potential benefit of AI, since it may bring breast cancer detection at mammography screening to the same high level everywhere.


133




134




135




136




137




138




139



140



141



142




143



144


# Aberrations of Normal Development and Involution ("ANDI")

- 1) Excess amount of fluid produced
  - a) Fibrocystic change, cysts




3D image of apocrine metaplasia

145




146




Multiple cysts surrounded by fibrosis

147



Comparative 3D histologic-galactographic images in fibrocystic change


148




149



150




151



152

*Fibrocystic change*




153

*Various sizes of cysts in fibrocystic change*




154




155

3D histology correlated with  
galactography and  
hand-held ultrasound




156

*Fibrocystic change*




157



158



159




160

## Demonstration of

- Normal ducts
- Dilated, fluid-filled ducts, ductectasia
- Cancer of ductal origin (DAB)


161

## 12-18 lobes



The 12-18 breast lobes have their own individual openings on the nipple surface. Preparation of Cooper (a) and thick section histology image in cross section showing the ducts within and behind the nipple (b).

162




*The major ducts are pleated. There is a slight twist on the pleated pattern.*

163




The pleats follow the twists in the major ducts.

164




165



166




167




168




169



170




171




172

Breast cysts


173




174



175




176




177



178



179



180

*Nipple discharge of the breast.*



Bologna, Italy

181



*Lack of indication for galactography*




Milky or turbid / cloudy discharge from multiple openings on the nipple.

182



183

*Indication for galactography*



*Bloody nipple discharge*

Cancer is the underlying cause:  
**13%**

184

### Indication for galactography




Serous nipple discharge




Cancer is the underlying cause: 7%

185




186

The procedure of choice is: *Galactography*  
Benign, intraductal papilloma cases




Histology: Benign papilloma without associated atypia.


187



188



189




190




Micropapillary fluid-producing carcinoma (DAB) in a major duct

191



Micropapillary ductal adenocarcinoma (DAB) producing fluid, which distends the duct

192




193

45-year-old woman, felt a thickening in the lower-lateral portion of her right breast. Also, she had bloody nipple discharge from the right nipple.


From the Departments of Mammography  
Surgery and Clinical Pathology  
Falun Central Hospital, Sweden ©

This slide features a blue text box at the top with clinical information. Below it is a photograph of a blue butterfly perched on a yellow hibiscus flower. At the bottom right, there is a small blue box with white text crediting the source to Falun Central Hospital, Sweden.

194




195




196




197




198



199



200




201




202




203



204




205



206



207



208



209